

Activities of Daily Living After Your Heart Procedure

Important precautions and guidelines

This handout gives important guidelines to follow to help your healing and keep you safe after your procedure.

Your Procedure	
You recently had this procedure:	
☐ Transfemoral valve replacement	3-4
☐ Transapical valve replacement	
☐ Mitral clip procedure	
☐ Cardiac catheterization	65 Po
<i>Type:</i>	
General Precautions	

Follow these precautions for:

How long:



After your procedure, closely follow the precautions in this handout.

For Your Safety

- Do **not** lift anything that weighs more than 5 to 10 pounds (a gallon of milk weighs almost 9 pounds). This includes grocery bags, laundry baskets, babies, and pets. Ask for help as needed.
- Do **not** hold your breath or strain or bear down while getting up or having a bowel movement.

Bathing

- You may shower. Gently clean your incision with soap and water and gently pat to dry. Do **not** rub or scrub.
- Do not soak in a bath tub or hot tub or go swimming.
- Do not apply any lotion or powder on your incision site.
- If it is hard for you to shower, your occupational therapist (OT) may advise you to use tools such as a shower chair, a hand-held shower head, a non-skid mat, a long-handled sponge, or a grab bar.

Getting Dressed

- To keep from falling, sit down to get dressed.
- If it is hard for you to put on your underwear, pants, socks, and shoes, your OT may advise you to use tools such as a reacher, dressing stick, sock aid, and long-handled shoehorn.

Extra Instructions

☐ Transapical Valve Replacement

If you had a transapical valve replacement, follow all instructions listed under "General Precautions."

Also follow these guidelines for 2 weeks when using your arm that is on the same side as your surgery (your surgical arm):

- Do **not** lift your elbow above your shoulder.
- Do **not** reach behind you when bathing, toileting, or putting on or taking off your clothes. Use your non-surgical arm.

□ Radial Artery Access

If you had your radial artery accessed for your procedure, follow all instructions listed under "General Precautions."

Also follow these guidelines when using your arm that is on the same side as your surgery (your surgical arm):

- For 48 hours, do **not** lift, push, or pull anything.
- For 5 days, do **not** lift anything that weighs more than 5 pounds. (A 2-liter bottle of water weighs almost 5 pounds.)

Questions?

Your questions are important. Call your doctor or healthcare provider if you have questions or concerns.

Occupational Therapy: 206.598.4830

5-Southeast CCU: 206.598.6500