UW Medicine

After Your Immunotherapy

Self-care and warning signs

Follow-up Visits

Bring your list of discharge medicines with you to your first follow-up visit at the clinic.

When to Call

Call the clinic if you have questions or any of the symptoms listed below. Please also see the "Quick Symptoms Guide" that starts on page 2.

Symptoms

- If you take your temperature by mouth:
 - Fever between 38 to 38.2° C (100.4 to 100.8° F) for 1 hour or more
 - Any fever 38.3°C (100.9°F) or higher
- If you take your temperature under your arm:
 - Fever between 37.5 to 37.7°C (99.5 to 99.9°F) for 1 hour or more
 - Any fever 37.8°C (100°F) or higher
- Chills or shaking, with or without a fever
- Signs of a cold, such as runny nose, watery eyes, sneezing, or coughing
- Changes in how your central line site looks, such as redness, swelling, pain, or drainage
- Changes in the color or consistency of your stool or urine
- New cough or shortness of breath
- Nausea, vomiting, or diarrhea
- Cannot take the medicine your provider prescribed for you
- New or different pain
- Any bleeding or bruising
- Headache or dizziness
- Change in your ability to think clearly



Call the clinic if you have any questions or any of the symptoms listed on this page.

Staying Healthy

Besides washing your hands, here are some other things that will help you stay healthy:

- Follow the line care instructions you received. Be sure to cover your line when you bathe or shower.
- Take your temperature 2 times a day, in the morning and at night.
- Avoid crowds and being around people who have cold or flu symptoms.

IV Therapy and Central Line Care

Your IV therapy and central line care are provided by:

- □ Seattle Cancer Care Alliance: 206.606.6000
- Derividence: 425.687.4400
- **C**oram: 425.883.3525

Quick Symptoms Guide

Here is a quick guide of warning signs to watch for, and what to do. Be sure to give a copy of this guide to your family members and caregivers.

	Emergency! Call 911 RIGHT AWAY If You Have These Symptoms Tell medics to take patient to UW Medical Center	Call Clinic NOW If You Have These Symptoms	
Areas of Concern		206.606.6000 Weekdays, 8 a.m. to 10 p.m. Weekends, 8 a.m. to 6 p.m. Holidays, 8 a.m. to 5 p.m.	After hours: 206.598.8902
Alertness, Consciousness, Activity	 Unconscious Cannot arouse Seizure Sudden change in vision Numbness, tingling, or cannot move arms or legs as usual One side of the face is drooping Cannot speak 	 New or increased confusion Change in level of alertness Mood changes, feeling irritable, f Change in ability to walk Trouble finding words Trouble swallowing Tremors or shakiness Dizziness Weak or <i>lethargic</i> (very tired, hat Trouble moving around 	

- □ Other:

Concerns	Call 911 Right Away	Call Clinic Now
Bleeding	Bleeding will not stop	New or increased bleeding
	Unconscious	Bloody urine
		New bruising
		Little red or purple spots on the skin
		Cannot stop nosebleed
		Bloody diarrhea
		Vomiting blood
		Using 1 or more feminine pads an hour
Blood Sugar	• Blood sugar lower than 50 and cannot waken	High or low blood sugar
		• Hard to wake up (use glucagon)
Breathing	Not breathing	Trouble breathing
8	• Choking or not moving	• Gets "winded" more easily with very little activity
	air into chest	• Feeling as if can't get enough air
		Trouble breathing when lying flat
		Wheezing with breaths
		New or recurring cough
		Persistent, continuous cough
		 Coughing blood or green-yellow material
Central	• Line open to air,	Line broken or leaking
Line (Hickman Line)	patient short of breath (Clamp line right away)	• Swelling in your face, neck, or exit site
		Cannot flush the line
		Line fell out
		Headache related to infusions
		• New redness, swelling, or tenderness at the exit site
		Drainage from exit site
Diarrhea		Constant or uncontrolled diarrhea
		New onset diarrhea
		Diarrhea with fever and abdominal cramping
		Whole pills passed in stool
		• Diarrhea more than 5 times a day
		Stool is bloody, deep red, or black
		Abdominal cramping
Fall	• Hit head during fall and change in	Did not hit head during fall, no changes in consciousness
		Report all falls to your provider
	consciousness after fall	
Fatigue	Cannot wake up	• Dizziness
_		Fatigue is getting worse
		• Too tired to get out of bed or walk to the bathroom
1		• Staying in bed all day

Concerns	Call 911 Right Away	Call Clinic Now	
Fever, Chills		 Chills or shaking chills, even if temperature is normal Fever (taken by mouth) between 38 to 38.2°C (100.4 to 100.8°F) for 1 hour or more, or a fever that is 38.3°C (100.9°F) or higher If taking steroids: Fever higher than 1 degree above usual Cold symptoms: runny nose, watery eyes, sneezing, coughing 	
Mouth Pain, Mucositis	 Not breathing Severe problems breathing 	 Trouble breathing Bright red blood in your mouth Pain not controlled by medicine White patches or sores appear on your gums or mouth Problems swallowing food or fluid 	
Nausea, Vomiting	• Vomiting that causes choking or problems breathing	 Nausea not controlled with anti-nausea medicines <i>Projectile</i> (explosive) vomiting Uncontrolled, constant nausea and vomiting Blood or "coffee ground" material in your vomit Can't keep medicine down because of vomiting Weakness or dizziness with nausea or vomiting Severe stomach pain while vomiting 	
Pain	 Severe chest or arm pain Severe squeezing or pressure in your chest Severe sudden headache 	 New or uncontrolled pain New headache Chest discomfort Heart pounding or feels like it's doing "flip-flops" Heart feels like it is "racing" Burning in your chest or stomach Pain or burning while urinating Painful central line site or in the area of the "tunnel" Pain with infusion of medicines or fluids into the central line 	
Rash		New rashRash with pain and or itching	
Swelling	• Throat swollen, feeling like you cannot get a breath	Sudden swelling, with or without painSwollen legs, arms, and hands	
Urination		 Cannot urinate for more than 8 hours Bloody urine Pain or burning while urinating 	

Questions?

Your questions are important. Call your doctor or healthcare provider if you have questions or concerns.