

## At Home After Your TAVR

### *Self-care and what to expect*


*This handout explains what to expect after your **transcatheter aortic valve replacement (TAVR)** and how to care for yourself at home. It includes pain control, incision care, exercising safely, and more.*

### Plan Ahead for Your Return Home

A successful TAVR means that you return home to live a better, longer life. Planning ahead is the most important thing you and your family can do to help make your TAVR a success.

**You will need help when you first go home.** Start planning for your return home as soon as you know you will be having TAVR.

- We suggest that a responsible adult stay with you for **at least 1 week** after you return home.
- It is hard to know how much help you will need or how long you will need it. Coping with your heart problem may have affected your overall health. You may need 1 to 2 months to fully recover after a successful TAVR.
- If you are not sure you will have the help and support you need at home, please ask to talk with one of our social workers. You may need home healthcare or even a skilled nursing or rehabilitation facility after TAVR.



*Plan ahead so you are sure you have the help you need at home after your TAVR.*

### Care Timeline

#### The First 30 Days after TAVR

The UWMC TAVR team will manage your care for the first 30 days after your procedure. After you go home, our team will monitor you closely to make sure your recovery is going well so that you do not need to be readmitted to the hospital.

- **In the first few days after discharge:** A cardiology nurse will call you to follow up on your symptoms, medicines, and well-being.
- **30 days after TAVR:** You will have a follow-up visit with a cardiology nurse practitioner or physician's assistant. At this visit, you will have an *echocardiogram* (ultrasound of the heart) to check how well your valve is working. You will also do a walk test and fill out a questionnaire about your symptoms and quality of life.

## 6 Weeks after TAVR

Usually 6 weeks after TAVR, our team will transfer your care back to your primary care provider and primary cardiologist. When this occurs, follow up with these providers.

Keep in mind that you may need to see your usual local healthcare providers sooner than 6 weeks.

## 1 Year after TAVR

You will see a member of the Structural Heart Team 1 year after your procedure. This follow-up visit will include an echocardiogram, a walk test, and a symptom/quality of life questionnaire.

## Research Study Participants

If you are in a research study, you may have follow-up visits more often.

## Dental Work After Your Procedure

Tell your dentist that you had a TAVR procedure. For 6 months after the TAVR procedure, you **must** have antibiotics before any dental work, even a dental exam. This is to prevent infection in your valve.

Your primary cardiologist can prescribe the antibiotics for you. If your dental office has any questions, please tell them to contact us.

## Aortic Stenosis and Congestive Heart Failure

People with aortic stenosis often have *congestive heart failure*, which is when your heart does not pump blood as well as it should. This occurs because the heart muscle has to work hard to pump blood through the narrow aortic valve, which makes the muscle thick and stiff.

Some symptoms of congestive heart failure are:

- Ankle and leg swelling
- Shortness of breath
- Fatigue

After TAVR, these changes in your heart muscle will take time to resolve. Your medicines may be changed to help ease symptoms.

## Symptoms of Congestive Heart Failure and What to Do

### OK



**I am OK if I can do my normal activities without a problem:**

- No shortness of breath
- Weight is stable
- No new swelling
- Normal appetite

### Warning



**I need to call my healthcare provider if my symptoms are getting worse:**

- Gained 3 pounds in 1 day or 5 pounds in 2 days
- New or increased swelling in my legs or ankles
- Harder time breathing, new cough, or I need to use more pillows to breathe while sleeping
- Nausea that will not go away, or I cannot eat
- Fatigue that is getting worse
- Increased heart rate (*palpitations*)

### Emergency!



**I need to call 911 right away if my symptoms are very bad or get worse quickly:**

- Chest pain that does not go away
- Severe dizzy spells or have fainted
- Severe or sudden shortness of breath
- New confusion or I cannot think clearly

## Your Fluid Balance

Congestive heart failure can cause you to have too much fluid in your body after TAVR. We manage your fluid balance with *diuretics* (water pills).

Weight gain is one sign of having too much fluid buildup in your body. At home, you will weigh yourself every day and keep a record of your weight. You received a handout called “Heart Failure Daily Weight Log” to record your weight and vital signs before you left the hospital. It is also attached to this handout. Please bring this log to your follow-up visits.

If you gain more than 2 pounds over what you weighed when you went home from the hospital, or if you have symptoms of congestive heart failure (see symptoms on page 2), call 206.598.6190 and ask to page the Cardiac Surgery Nurse or Nurse Practitioner on call.

Even though you may have extra fluid in your body, you still must stay hydrated. The right amount of liquid to have each day will vary, but most people need to drink about 1 to 2 liters of liquid a day.

If you are retaining more fluid or having more symptoms, you may be placed on a *fluid-restricted diet*. We will review this with you in follow-up phone calls and office visits.

## Managing Your Pain and Changes in Sensations

You can expect to have some aches and pains after TAVR. You may have pain in your back, shoulders, neck, and chest. These are part of the normal healing process and may last up to a month.

There may also be changes in how your body feels heat and cold, how your body regulates your temperature, and how things taste. These symptoms will go away over time.

You may leave the hospital with a prescription for pain medicine. If you are sore or uncomfortable, please take this medicine as directed. If you do not take your pain medicine, you may not be as active as you should be.

If your pain medicine is not working well, call 206.598.6190. Ask to page the cardiac surgery nurse or nurse practitioner on call.

## Showering and Bathing

- You may take a warm shower when you get home, unless your doctor tells you not to. When you shower:
  - Gently wash your incisions with mild, unscented soap and water.
  - Pat your incisions dry (do not rub them).
- You may take a warm bath after your incisions heal fully, about 2 weeks after surgery. Do **not** soak your incisions for longer than 5 minutes until all the scabs have fallen off and your incisions are fully closed.
- Do **not** take very hot showers or baths or soak in a hot tub. Hot water may lower your blood pressure. This could make you weak or dizzy or cause you to faint. (Medicines you are taking may also cause fainting.)

## Care of Your Incision and Wrist Puncture Site

### Signs of Infection

Watch your incision closely for any signs of infection. These include:

- Redness, warmth, or heat around the incision
- New drainage
- Fever of 101°F (38.5°C) or higher
- Bleeding at the wrist or groin site that does not stop after applying pressure for 10 minutes

If you have any of these signs, call one of these numbers **right away**:

- During clinic hours, call **206.598.VALV (8258)** and ask for the nurse or nurse practitioner.
- After hours and on holidays and weekends, call **206.598.6190** and ask to page the CARD I Structural Fellow on call.

### Incision Care

Most patients have a bruise at the incision site on their groin. This bruise may spread down your leg over the next day. It can take 2 to 3 weeks to go away.

- If your incisions are:
  - **Still draining** when you leave the hospital, your nurse will give you instructions and supplies to care for them at home.
  - **Not draining**, leave them open to the air. If your clothing rubs or irritates them, you can cover the area with dry gauze. Remove the gauze at night. You do not need to apply any special dressing.
- Do **not** put any medicine, lotion, or powder on your incisions until they are fully healed and the scabs have fallen off, unless your care team tells you otherwise.

### Wrist Puncture Site

- Do **not** allow the puncture site to be covered by water. This means do not take a bath, sit in a hot tub, or go swimming. It is OK to take a shower.
- You may remove the wrist dressing (bandage) the day after your procedure.
- It is normal to have a small bruise or lump at the puncture site.

## Leg Swelling

Some people have a lot of swelling in their legs after TAVR surgery. If you do, we may advise you to wear *antiembolic* stockings. These stockings lessen swelling, improve blood flow, and help keep blood clots from forming in your legs.

- For 3 weeks after your surgery, wear the stockings when you are out of bed. Take them off at night.
- After 3 weeks, wear them only if you still have swelling in your legs.
- If you get new or increased swelling, see “Warning” section on page 3.

## Activities

After you get home, you can expect to slowly return to a normal lifestyle. But, be careful not to do things that make your heart work too hard.

How active you are and what type of activities you can do after you leave the hospital depend on:

- Your condition before surgery
- The type of surgery you had
- How your recovery is going

As soon as you wish, you may:

- Shower, shave, and wash your hair.
- Walk on level ground at an easy pace. Slow down when you walk up hills. (See “Your Walking Program” on page 8.)
- Walk up and down stairs at a normal pace, but limit how much you do this. Remember that going up takes more energy. Slow down or stop when you become short of breath.
- Use a stationary bicycle.
- Ride in a car, go out for meals, or visit friends.
- Prepare meals.
- Wash dishes and clothes (see precautions below).
- Do light housekeeping, such as dusting (see precautions below).
- Resume sexual activity (see pages 10 and 11).

## Activity Precautions After Your Procedure

### Protect Your Wrist

<b>For 48 hours:</b>	<ul style="list-style-type: none"><li>• <b>AVOID</b> lifting, pushing or pulling with the affected arm.</li><li>• <b>AVOID</b> bending, turning, or twisting your wrist on the affected arm.</li><li>• Do <b>NOT</b> have your blood pressure taken on the affected arm.</li></ul>
<b>For 5 days:</b>	<ul style="list-style-type: none"><li>• Avoid vigorous exercise that uses the affected arm.</li><li>• Do <b>NOT</b> lift anything that weighs more than 5 pounds (2.27 kilograms) with the affected arm. (A 2-liter soda bottle weighs more than 4 pounds.)</li><li>• It is normal to have a small bruise or lump at the insertion site.</li></ul>

### Protect Your Groin

You may have a bruise at the insertion site on your groin. This is normal. It may spread down your leg over the next day. It may take **2 to 3 weeks** to go away.

- For **7 days** after TAVR, do **NOT**:
  - Lift anything that weighs more than 10 pounds (4.54 kilograms). This includes pets, groceries, children, trash, and laundry. (A gallon of milk weighs almost 9 pounds.)
  - Garden, including mowing the lawn and raking.
  - Hold your breath, bear down, or strain when having a bowel movement.
  - Allow the puncture site to be covered by water. This means do **not** take a bath, sit in a hot tub, or go swimming.
- For **30 days** after TAVR (until you have your 30-day follow-up visit), do **not**:
  - Run, jog, swim, or bike
  - Play sports like golf, tennis, bowling, or softball

### If You Had Transapical or Transaxillary TAVR

For **2 weeks** after TAVR:

- Do **NOT** lift, push, pull, or carry anything that weighs more than 10 pounds (4.54 kilograms). This includes groceries, children, pets, trash, and laundry. (A gallon of milk weighs almost 9 pounds.)



*When you go walking, wear comfortable, flat shoes and loose clothing.*

- Do **NOT** reach your arms behind your back or above shoulder level. Your goal is to keep your skin and muscle from stretching. This will help your incision heal faster.
- Do **NOT** mow the lawn, rake, or do other work in the garden.
- Do **NOT** hold your breath, bear down, or strain when having a bowel movement.

## Driving

- Do **not** drive for 1 to 4 weeks after TAVR.
- Some people have other medical reasons to restrict driving. If this is true for you, your primary doctor(s) may need to make the decision about when you can drive.

## Exercise Safely

Daily exercise is vital to healing. Regular exercise may strengthen your heart muscle and allow it to pump blood better. Exercise also improves muscle tone and blood flow in your body, helps with weight loss or maintaining your weight, and may create a general sense of well-being.

When you exercise:

- If your doctor or physical therapist advises you to use an assistive device such as a cane or walker, **be sure to use the right device needed to keep you safe and prevent falls.** It is important to stay active, but staying safe is more important than walking on your own without a cane or walker.
- Wait 1 hour after eating before you walk or do any other exercise.
- Wear comfortable, flat shoes and loose clothing. Any clothes that restrict your movements may affect your breathing.
- Stretch your muscles before and after exercising. This will help you get the most from your exercise and will help prevent injuries.
- Take your pulse on the unaffected arm to see how hard your heart is working. **Do not let your pulse get higher than 20 to 30 beats a minute above your resting heart rate.** (See “How to Take Your Pulse” on page 10.)
- When you are walking long distances, stay on level ground. Going up hills and stairs is OK if you go slowly and keep your heart rate within your safe guidelines.
- Avoid being outside on very hot or very cold days during your recovery. Very high and very low temperatures stress your heart.

In the winter, walk in the afternoon or during the warmest part of the day. During the summer, walk in the morning or during the coolest part of the day.

### **Your Walking Program**

How long and how far you walk will depend on how well you are handling the exercise. Remember to use your pulse as your guide to tell you how hard your heart is working (see page 10).

As you get stronger, you will begin to walk a little farther and a little faster. At first, increase the distance you walk before increasing how fast you walk.

### **Slow-down Signals**

Your body will tell you if the exercise you are doing is too hard for you. Slow down or stop what you are doing, write down what happened, and talk with your nurse or doctor if:

- Your exercise heart rate or pulse is more than 20 to 30 beats above your resting heart rate.
- You cannot talk while exercising. Slow your activity until you have control of your breathing and can talk without being breathless.
- Your heart rate or pulse stays high for 10 minutes after you stop exercising.
- You cannot catch your breath for longer than 10 minutes after you stop exercising.
- You are tired for up to 24 hours after exercising.
- You have pain in your joints, shins, or heels.
- You have pain or cramping in your calf muscles.

### **When to Stop Exercising**

Stop exercising and talk with your doctor before you start your exercise program again if you:

- Have an abnormal heart rhythm, such as:
  - An erratic pulse or palpitations
  - Your pulse suddenly gets very slow
  - A sudden burst of rapid heartbeats
- Have new pain, pain that lasts a long time, or pressure in your chest, arms, or throat
- Are dizzy, confused, or lightheaded
- Lose control of your muscles or faint

- Have cold sweats or become pale
- Feel sick to your stomach or are vomiting

Do **not** exercise if you have:

- A bad cold, flu, or fever
- Fatigue (feel very tired)

### Warm-up Exercises

Warm-up exercises help your body get ready for activity. Do your warm-up exercises 2 times a day – once in the morning and once in the afternoon. Do them slowly and steadily, and keep breathing slowly (do not hold your breath).

Warm-up exercises include:

- Marching in place
- Leg stretches (if you learned these in physical therapy after TAVR)

Begin by doing each warm-up exercise 5 times. Each time you do an exercise is called a *repetition* or “rep.” Over time, you can increase the number of reps, but do not do more than 20 reps at a time.

Do **not** add more reps if you have any of the symptoms listed in the “Slow-down Signals” section on page 8 or in the “Warning Signs” section above.

### How to Take Your Pulse

Your best guide to how hard your heart is working is your pulse (heart rate). When you feel your pulse, you are actually feeling your heart pushing blood through your arteries.

Do **not** take your pulse on your affected arm. Check it on the other wrist.

### Steps

To take your pulse:

- Use your fingers (**not** your thumb, which has a pulse of its own) to find your pulse on the inner part of your other wrist, just below your thumb.
  - If you cannot find your wrist pulse, gently find the pulse in your neck. Do not press hard. You could reduce blood flow to your head, making you feel dizzy or faint.
- Start with 0, and count your pulse for 1 minute. You can also count your pulse for 15 seconds and multiply that number by 4.



Take your pulse on the wrist of your **unaffected arm**.

## When to Take Your Pulse

Check your pulse:

- When you are at rest
- During exercise
- At the end of exercise
- If you have symptoms such as hard or thumping heartbeats, *palpitations* (rapid, fluttering heartbeats), dizziness, or shortness of breath that do not go away when you rest

Remember: Do not let your pulse get higher than 20 to 30 beats a minute above your resting heart rate.

## Resuming Sexual Activity

Wait at least 2 weeks before you resume sexual activity. If you have any questions or concerns after that time, please talk with your healthcare provider.

## Returning to Work

When you can return to work will depend on your recovery and what kind of work you do. Talk about your return to work with your healthcare provider at your 30-day follow-up visit. Most patients are able to return to full-time work within 3 months after surgery. Many return earlier.

## Becoming More Independent After Surgery

Before surgery, you may have become more dependent on loved ones as your health got worse. After successful heart surgery, you will start becoming more independent and need less help from others. For your safety and well-being, you and your loved ones will have to decide how to balance your independence with having help when you need it.

## Warning Signs

These symptoms are **warning signs**:

- Chest pain when you are active
- Shortness of breath that does not improve with rest
- *Palpitations*, fluttering in the chest, or a hard, thumping heart beat
- Dizziness, feeling faint, or feeling like you may pass out
- Unexpected, sudden changes in your vision, strength, or coordination

If you have any of the warning signs listed above:

- Sit down or lie down and rest.
- Then take your blood pressure and pulse.
- If your symptoms do not improve after a few minutes of rest, call 206.598.6190 and ask to page the CARD I Structural Fellow on call.

If you have any urgent concerns, call 911.

## If You Have More Questions

It is normal to feel anxious about going home after your TAVR surgery. If you have questions or concerns about any issues that come up, please call us.

### Questions?

Your questions are important. Call your doctor or healthcare provider if you have questions or concerns.

**Heart Institute at UWMC:** Call 206.598.VALV (206.598.8258) during clinic hours and ask for the nurse or nurse practitioner.

**After clinic hours or on holidays or weekends:** Call 206.598.6190 and ask to page the CARD I Structural Fellow on call.