

# **Exparel Injections**

Bupivacaine liposomal

This handout explains a medicine called Exparel, a local anesthetic that is used to prevent or treat pain from some procedures.

## What is Exparel?

Exparel is the brand name for a drug called *bupivacaine liposomal* (pronounced bue-PIV-a-kane LIP-oh-som-al). This drug is a *local anesthetic* (numbing medicine). It causes loss of feeling in the skin or other tissues.

### How is it used?

Exparel is used to numb an area of your body before or after minor procedures. It is given by injection by a healthcare provider in a hospital or clinic.

If this medicine will be given to a child, talk with your pediatrician first. Special care may be needed.

## How do I prepare for this injection?

Tell your provider if you have:

- Heart disease
- · Kidney disease
- Liver disease
- An unusual or allergic reaction to bupivacaine, other medicines, foods, dyes, or preservatives

#### Women

Tell your provider if you are:

- Pregnant or trying to get pregnant
- Breastfeeding



Talk with your nurse or doctor if you have any questions about Exparel.

## **Does Exparel have side effects?**

Tell your doctor or other provider **right away** if you have any of these side effects after your Exparel injection:

- Allergic reactions such as:
  - A rash, itching or hives
  - Swelling of your face, lips, or tongue
- Breathing problems
- · Changes in vision
- Feeling dizzy
- · Fast or slow, irregular heartbeat
- Joint pain, stiffness, or loss of motion
- Seizures

These side effects can be bothersome, but do not usually require a visit to the doctor:

- Constipation
- · Irritation at site where injected
- Nausea or vomiting
- Feeling very tired

Call your provider if any of your symptoms are causing you concern.

### What else do I need to know?

- Your condition will be monitored carefully while you are receiving Exparel.
- Be careful to avoid injury while the injected area is numb. You will not feel pain, so you can easily hurt yourself.
- For 96 hours after receiving Exparel:
  - Do not use another medicine that contains bupivacaine. You may have more side effects.
  - Do not have other anesthetics.

Talk with your doctor if you have any questions or concerns.

### **Questions?**

Your questions are important. Talk with your doctor or other healthcare provider if you have questions or concerns.