UW Medicine

UNIVERSITY OF WASHINGTON MEDICAL CENTER

Mastectomy with Axillary Node Dissection CareMap

Before, during, and after your hospital stay

Before Surgery	Day of Surgery	Discharge or Day 1
7 Days Before ☐ Plan for your ride to the hospital and back home. ☐ If you usually take blood thinners such as Lovenox or Coumadin, talk with our care team. We may want to adjust your doses around the time of the surgery. ☐ Stop taking supplements and vitamins. ☐ Pre-Anesthesia staff will tell you which medicines to take on the morning of surgery. 2 Days Before ☐ Stop shaving near the surgery site, if needed. Day Before ☐ You will receive a call from the hospital with your check-in time and medicine instructions. ☐ Take a shower with the antibacterial soap, as prescribed. ☐ Do NOT eat any food or drink any alcohol after midnight. You may drink cliear liquids up until 2 hours before your check-in time. ☐ Read the section on	Before Surgery At Home Take a 2nd shower with the antibacterial soap, as prescribed. At the Hospital Check in at the Surgery Registration Desk at or before your check-in time. Anurse will call you to come into the pre-op area. We will start an intravenous (IV) line to give you medicines and fluid. We may check your blood sugar. You will meet with: An anesthesiologist to talk about anesthesia A member of the surgery team so that you can ask questions and sign a consent form (if not already signed) Nurses to review your health history A member of the anesthesia team will take you to the operating room (OR). After Surgery You will wake up in the recovery room. You will: Have drains to remove fluid from the surgery area Receive pain medicine by IV or by mouth, and anti-nausea medicine as needed Self-care and Activity Arm-use precautions: Until your drains are removed, move your arms gently. Do NOT raise your arm above shoulder height on the side of your surgery. Do NOT lift anything that weighs more than 8 pounds (1 gallon of water weighs more than 8 pounds). This includes children and pets. Do NOT vacuum, do laundry, or do other chores. Do not use the arm on the side of your surgery for blood pressure checks, IV placements, or blood draws.	 □ Once your pain is under control, you will be discharged from the hospital. This could be the same day as surgery or the next morning. □ Your follow-up visit with your surgical team will be set up before you leave the hospital. Medicines □ Your pain will be controlled by: Acetaminophen (Tylenol) and/or ibuprofen (Advil, Motrin) Opioid pain pills, only if other medicines to do ease your pain (always take with food) □ While you are taking opioids, also take the medicine prescribed for constipation (unless you have loose stools). Diet □ Eat your usual healthy foods. □ Drink plenty of water to stay hydrated. Self-care and Activity □ If you have dressings, leave them in place. □ Keep the drain site clean. If you wish, you may lightly cover the site with gauze and tape. For more deails, read the "Drain Care" section of your manual, Guide to Your Breast Surgery. □ Empty each drain 2 to 3 times a day. Log each amount separately. Call the clinic when output is less than 30 ml in 24 hours for 2 days in a row. They will tell you next steps for drain removal. □ Walk at least 4 times a day. Start with short walks, about 10 to 15 minutes long. □ Do NOT drive while you are taking opioids. □ Read and follow instructions in the handout "Stretches and Exercises." Move your arms gently. Follow arm-use precautions until drains are removed (see "Self-care and Activity" in Day of Surgery column). At Home □ Continue all medicines, diet, self-care, and activity instructions
Read the section on preparing for surgery in your manual Guide to Your Breast Surgery.		 Continue all medicines, diet, self-care, and activity instructions provided at discharge. 48 hours after surgery: If you have dressings, remove the outer bandages from your breast and armpit. Leave the white Steristrips in place. Then shower. Pat the Steri-strips dry.

After Drains Removed Week 1 Weeks 2-3 **Medicines Medicines** Medicines ☐ Pain controlled by: ☐ Pain controlled by acetaminophen and ibuprofen ☐ Pain controlled by acetaminophen and ibuprofen - Acetaminophen and ibuprofen ☐ Goal is to be off opioids by now **Self-care and Activity** - Opioid pain pills, only as needed (always take with food) **Self-care and Activity** ☐ Walk every day, going farther ☐ Walk every day, going farther each day. ☐ Begin to taper dose of opioids each day. ☐ While you are taking opioids, keep taking the medicine prescribed ☐ Arm-use precautions: Until drains are ☐ Until you see a physical or for constipation (unless you have loose stools). removed, move your arms gently: occupational therapist (PT or OT), - Do **NOT** raise your arm above shoulder **Self-care and Activity** keep following instructions in the Walking height on the side of your surgery. Activity "Stretches and Exercises" handout. ☐ Walk every day, going farther each day. Guidelines - Do **NOT** lift anything that weighs more ☐ Do **NOT** use the arm on the side ☐ Arm-use precautions: Until drains are removed, than 8 pounds (1 gallon of water weighs of your surgery for blood pressure move your arms gently: almost 9 pounds). This includes children checks, IV placements, or blood - Do **NOT** raise your arm above shoulder height and pets. draws. on the side of your surgery. Do **NOT** vacuum, do laundry, or do other chores. When your surgeon says it is safe: - Do **NOT** lift anything that weighs more than 8 pounds (1 gallon of - Do **NOT** use the arm on the side of your surgery for blood ☐ You may start PT/OT and massage water weighs almost 9 pounds). This includes children and pets. pressure checks, IV placements, or blood draws. 1 week after your last drain is - Do **NOT** vacuum, do laundry, or do other chores. removed. ☐ Keep following instructions in the "Stretches and Exercises" - Do **NOT** use the arm on the side of your surgery for blood handout. ☐ Start using your arms more fully. pressure checks, IV placements, or blood draws. Drop weight limits. ☐ If you still have drains, empty each of them 2 to 3 times a ☐ Keep following instructions in the "Stretches and Exercises" day, record each amount separately in your log. Call the ☐ Start doing more exercise. Build handout. clinic to have a drain removed when output for that drain is slowly. Let your body guide you. less than 30 ml in 24 hours for 2 days in a row. If an activity causes pain, slow ☐ Empty each drain 2 to 3 times a day. Record each drain amount down or stop! separately in your log. Always bring your log with you to your clinic ☐ If you have stopped taking opioids, you can drive if you feel visit. OK doing so and you follow your activity restrictions. ☐ Do **NOT** drive while taking opioids. **Follow-up Visits Ouestions or Concerns?** ☐ Visit with your breast surgeon and/or Advanced Practice Practitioner (APP) 1 to 2 weeks after surgery: Follow-up Your questions are important. Call your care team if you have questions - Talk about pathology report (you may receive results by phone before this visit) or concerns. - Drains are removed, if output less than 30 ml a day (only 1 drain may be removed at your first follow-up visit) ☐ UWMC Breast Clinic - Northwest - If needed, talk about seeing medical or radiation oncologist for more treatment Campus: 206.668.6746 - Receive prescription for physical therapy (PT) or occupational therapy (OT), but do **not** use it until at least ☐ SCCA Breast Health Clinic: 1 week after all drains are removed 206.606.7563 For urgent needs after hours: You can also call your clinic any time

of the day or night. Ask to speak with

the provider on call.