UW Medicine

UNIVERSITY OF WASHINGTON MEDICAL CENTER

Mastectomy with Tissue Expander CareMap Before, during, and after your hospital stay

☐ Plan for your ride to ☐ Take a 2nd shower with the antibacterial soap, as prescribed. discharged from	s under control, you will be the hospital. This could be the gery or the next morning. isit with your surgical team will be
home. If you usually take blood thinners such as Lovenox or Coumadin, talk with your care team. We waill start an intravenous (IV) line to give you medicines and fluid. We will start an intravenous (IV) line to give you medicines and fluid. We will start an intravenous (IV) line to give you medicines and fluid. We will start an intravenous (IV) line to give you medicines and fluid. We will check your blood sugar. You will meet with: An anesthesiologist to talk about anesthesia the will take you to operating room (orange) While you are tall constitution of the morning of your surgery. A member of the anesthesia team will take you to operating room (orange) While you are tall constitution of the morning of your will wake up in the recovery room. You will wake up in the recovery room. You will wake up in the recovery room. You will wake up in the surgery area to remove fluid Fare as shower with the antibacterial soap, as prescribed. Do NOT eat any food or drink alcohol after midnight. Do NOT eat any food or drink alcohol after midnight. Walk starting the pight of surgery. Walk startin	controlled by: en (Tylenol) and ibuprofen (Advil, Motrin) ints els, if needed (always take with food) king opioids, also take the medicine prescribed for less you have loose stools). ealthy diet. er to stay hydrated. rity sings, leave them in place. ite clean. If you wish, you may lightly cover the site ape. Follow any other instructions you were given while you were in the hospital. imes a day. Start with short walks about 10 to

Week 1	Weeks 2 and 3	Week 4
Medicines	Medicines	Medicines
□ Pain controlled by:- Acetaminophen, ibuprofen, and muscle relaxants.	☐ Pain controlled by Acetaminophen, ibuprofen, and muscle relaxants as needed.	☐ Pain controlled by Acetaminophen, ibuprofen, and muscle relaxants as needed.
 Opioid pain pills, only as needed (always take with food). 	☐ Goal is to be off opioids by now.☐ Keep taking antibiotics, if you still have a drain in	☐ May need muscle relaxants for only 1 to 2 days after tissue expansion.
☐ Begin to taper your opioid dose.	place.	Self-care and Activity
☐ While you are taking opioids, keep taking the medicine prescribed for constipation (unless you have loose stools).	Self-care and Activity Walk every day, going farther each day. For 4 weeks, move your arms gently:	 □ Walk every day, going farther each day. □ For 4 weeks, move your arms gently: - Do NOT raise your arm above shoulder height on the
☐ Take antibiotics as prescribed to reduce infection risk.	 Do NOT raise your arm above shoulder height on the side of surgery. 	side of surgery. - Do NOT lift anything that weighs more than 8 pounds.
Self-care and Activity ☐ Walk every day, going farther each day.	- Do NOT lift anything that weighs more than 8 pounds. (1 gallon of water weighs more than	(1 gallon of water weighs more than 8 pounds.) This includes children and pets.
□ For 4 weeks, move your arms gently:Do NOT raise your arm above	8 pounds.) This includes children and pets. - Do NOT vacuum or do laundry or other chores.	 Do NOT vacuum or do laundry or other chores. Until your PT or OT visit, keep following instructions in the "Stretches and Exercises" handout.
shoulder height on the side of surgery.	☐ Keep following instructions in the "Stretches and Exercises" handout.	Week 5
 Do NOT lift anything that weighs more than 8 pounds. (1 gallon of water weighs more than 8 pounds.) This includes children and pets. 	☐ If you still have drains, empty them 2 to 3 times a day, record amounts in your log. Call clinic to have them removed when output is less than 20 ml in	Self-care and Activity When your surgeon says it is safe:
 Do NOT vacuum or do laundry or other chores. Keep following instructions in the "Stretches and Exercises" handout. 	24 hours for 2 days in a row. ☐ If drains are still in place at the end of 3 weeks, make an appointment to have them removed,	 You may start PT, OT, and massage. Start using your arms more fully. Drop weight limits. Start doing more exercise, build slowly, let your body
☐ Empty drains 2 to 3 times a day and record amounts in your log. Always bring the log with you	either in the breast surgeon's or plastic surgeon's office.	guide you – if it hurts, slow down or stop! Follow-up Visits
to your clinic visit.	If you have stopped taking opioids, you can drive	Visit plastic surgeon's office for tissue expansion:
☐ Do not drive while taking opioids. Follow-up Visits	if you feel OK doing so. Be sure to follow your arm use precautions.	☐ If your incisions are healed well enough, we will start your tissue expansion:
Visit with your breast surgeon or a Advanced Practice Practitioner (APP) 1 to 2 weeks after surgery:	Follow-up Visits Visit with your plastic surgeon or APP 3 weeks after	Every 2 weeks, orMore often if radiation is planned for shortly after
☐ Talk about your pathology report. (You may receive results by phone before this visit.)	surgery: If your incisions are healed well enough, we will	Surgery Questions or Concerns? Who to Call
☐ Drains are removed, if output less than 20 mL a day.	start your tissue expansion. We will remove any remaining drains, if output less than 20 mL a day.	Your questions are important. Call your care team if you have questions or concerns.
 ☐ If needed, talk about seeing a medical or radiation oncologist for more treatment. ☐ Receive a prescription for physical therapy (PT) or 	☐ A follow-up visit with plastic surgery will be set up by your Patient Care Coordinator. To confirm	 SCCA Breast Health Clinic: 206.606.7563 UWMC Breast Clinic Northwest Campus: 206.668.6746
occupational therapy (OT), but do not use it until 4 weeks after surgery.	or schedule your appointment, call the UWMC Plastic Surgery Clinic at 206.598.1217.	For urgent needs after hours: Call your clinic any time of the day or night. Ask to speak with the provider on call.